

CD10 (55C6)

Mouse monoclonal Antibody

Host: Mouse
Specificity: Human
Immunogen: Recombinant external domain of the human CD glycoprotein
Ig Class: IgG1
Storage: Store at 2-8°C for up to 2 years for concentrate form and 18 months for predilute form.

Specifications: CD10 (55C6) Mouse monoclonal primary antibody detects CD10 protein in formalin-fixed, paraffin-embedded samples by immunohistochemical (IHC) staining.

Staining procedures: Use formalin-fixed and paraffin-embedded sections. Retrieval conditions: Pretreatment of deparaffinized samples via heat-induced epitope retrieval is recommended. Detection method: Polymer anti-mouse/rabbit Ig detection system. Working dilution: 1:50-100. Positive Control: sample known to contain CD10 protein.

Localization: Membrane/cytoplasmic.

Intended Use: Research Use Only (RUO). This antibody is not for diagnostic use.

Description: The common acute lymphoblastic leukemia antigen (CALLA/CD10) is a single-pass type II transmembrane metallo-endopeptidase that cleaves and inactivates various peptide growth factors essential for signal transduction. CD10 is expressed by a few hematopoietic cells, such as immature T and B cells, B cells of the germinal centers of lymphoid follicles, and granulocytes. It also reacts with various non-hematopoietic cells, including epithelial cells in the GI tract and kidney tubules. In the liver, the bile canaliculi show a moderate to strong staining. CD10 is expressed in a high percentage of cases of acute lymphoblastic leukemia, follicular lymphoma, Burkitt's lymphoma, some hematopoietic tumors, and chronic myelogenous leukemias in lymphoid blast crisis. It is also known to be a marker of endometrial stromal cells. It helps differentiate endometrial stromal sarcoma (ESS) from uterine cellular leiomyoma (UCL) and uterine leiomyosarcoma (ULS).

Supplied As: Purified antibody in Tris-HCl pH 7.4 buffer containing stabilizing proteins (including < 1% Bovine Serum Albumin) and < 0.1% ProClin by volume.

References:

1. Maguer-Satta V, et al. Stem Cells. 2011; 29:389-96.
2. Onak NK, et al. Turk J Med Sci. 2010; 40:177-84.
3. Chu P, et al. Am J Clin Pathol. 2000; 113: 374-82.



Z2809ML-R/ Z2809MS-R/ Z2809MT-R/ Z2809MP-R (1.0ml Concentrate/ 0.5ml Con./ 0.1ml Con./ 7ml Pre-dil)



605 E. Huntington Dr.,
Suite 204
Monrovia, CA 91016
Tel: (626) 355-2053
<http://www.zeta-corp.com>



(research
use only)



(lot number)



(catalogue
number)



(expiry date)



(temperature
limit)



(consult
Instructions
for Use)